

WORLD INTELLECTUAL PROPERTY ORGANIZATION



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:

C07J 71/00

(11) International Publication Number:

WO 91/04984

A1

(43) International Publication Date:

18 April 1991 (18.04.91)

(21) International Application Number:

PCT/SE90/00619

(22) International Filing Date:

27 September 1990 (27.09.90)

(30) Priority data:

8903219-7

2 October 1989 (02.10.89)

SE

(71) Applicant (for all designated States except US): AKTIEBO-LAGET ASTRA [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and

(72) Inventors; and
(75) Inventors/Applicants (for US only): HOFSTRAAT, Robert,
Gerrit [NL/NL]; Zwanenveld 25-32, NL-6538 NK
Nijmegen (NL). RAIJMAKERS, Petrus, Hendricus
[NL/NL]; 11llostraat 19, NL-5402 AA Uden (NL).
VRIJHOF, Pieter [NL/NL]; Margrietlaan 10, NL-5351 BX Berghem (NL).

(74) Agents: MIKSCHE, Gerhard et al.; Aktiebolaget Astra, Patent Department, S-151 85 Södertälje (SE).

(81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CM (OAPI patent), DE (European patent)*, DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL (European patent), NO, RO, SD, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US.

Published

With international search report.

(54) Title: PROCESS FOR THE MANUFACTURE OF BUDESONIDE

(III)

(57) Abstract

The present invention relates to a novel process for the manufacture of (22 R,S)-16a, 17a-butylidenedioxy-11β, 21-dihydroxypregna-1,4-diene-3,20-dione (I) by reacting 11\(\beta\), 16\(\alpha\), 17\(\alpha\), 21-tetrahydroxypregna-1,4-diene-3,20-dione (II) with butanal, CH₃CH₂CH₂CHO in acetonitrile with p-tolyuenesulphonic acid as a catalyst.

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

ΑT	Austria	ES	Spain	MC	Monaco
AU	Australia	Fl	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Fasso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	PL	Poland
CA	Canada	JP	Japan	RO	Romania
CF	Central African Republic	KP	Democratic People's Republic	SD	Sudan
CG	Congo		of Korea	SE	Swedon
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CM	Cameroon	LI	Liechtenstein	SU	Soviet Union
DE	Germany	LK	Sri Lanka	TD	Chad
DK	Denmark	LU	Luxembourg	TG	Togo
_ /-			-	US	United States of America

Process for the manufacture of Budesonide

5 Technical field

The present invention relates to a novel process for the manufacture of (22 R,S)-16a,17a-butylidenedioxy-11B,21-dihydroxypregna-1,4-diene-3,20-dione (budesonide)


10 .

20

by reacting 118,16a,17a,21-tetrahydroxypregna-1,4-diene-3,20-dione (16a-hydroxyprednisolone)

25

30



35

with butanal, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$, in a solvent medium in the presence of an acid catalyst.

Prior art

According to a previously known process disclosed in GB patent no. 1 429 922 Budesonide is manufactured by reacting 16α-hydroxyprednisolone with butanal in dioxane and with perchloric acid as a catalyst. The product is recovered by diluting the re-5 action mixture with methylene chloride, and neutralising by washing with aqueous potassium carbonate and water, evaporating the solvent followed by crystallisation from ether/ligroine. The product was further purified by chromatography e.g. on Sephadex. The main disadvantages of dioxane are its skin penetrating and 10 peroxide formation properties. Another disadvantage with this prior art process is perchloric acid, which is a strong exidizing agent and the use of this catalyst results in a less selective reaction, which in turn makes the subsequent work-up and purification process complicated and expensive. 15

Disclosure of the invention

The object of the invention is to create a novel process, which gives a more selective reaction and a more simple and economic work-up and purification process.

This is achieved with the process according to the present invention, wherein the reaction is performed in acetonitrile with p-toluenesulphonic acid as a catalyst.

25

30

35

20

The combination of the less basic (compared to dioxane) solvent acetonitril and the weaker, i.e. non-oxidizing p-toluenesulphonic acid gives a more selective reaction, and also a more simple and exonomic work-up and purification process compared to the above discussed prior art process using dioxane and perchloric acid.

According to a preferred embodiment of the invention the reaction is stopped by the addition of water and adjustment of the pH of the reaction mixture. This might be done by the addition of sodium hydrogen carbonate in water. The product then crystallizes. The crystals are filtered off, dissolved in methylene chloride and methanol and are then crystallized by the addition

a suitable hydrocarbon, such as ligroine, hexane, cyclohexane or heptane, giving a crude product, which is then recrystallized in methanol/water to give pure budesonide.

5 The process according to the invention for the manufacture of budesonide thus consists of two steps.

Step 1. Budesonide crude

16α-hydroxyprednisolone is reacted with butanal in acetonitrile.

10 p.Toluenesulphonic acid is added as a catalyst. The reaction mixture is diluted with water and aqueous sodium hydrogen carbonate. After cooling to 5.15°C the crystallized product is filtered off and washed with water. The wet or dried substance is then dissolved in methylene chloride. If the substance used is wet the water phase formed upon dissolution is removed. Methanol is added and the resulting crude budesonide is precipitated by the addition of ligroine or another suitable hydrocarbon (e.g. hexane, heptane or cyclohexane) and is then filtered off.

20 Step 2. Budesonide

The crude budesonide is dissolved in methanol at about 60°C. The solution is filtered through a closed filter and the product is crystallized by the addition of water. After cooling to 5-20°C, filtration and washing with methanol/water the budesonide is dried in vacuum at 40-45°C.

This process is simplified, more economic and less health hazardous compared to prior art processes.

30 Working example

25

The reaction is carried out in a nitrogen atmosphere. 15,4 g ptoluenesulphonic acid is dissolved into 200 ml acetonitrile. To
the solution 50,0 g 16a-hydroxyprednisolone and 17.6 ml butanal
are added. The temperature rises to 25°C. After 30 min most of
the material is dissolved. Shortly thereafter the product starts
to crystallize. After 3 hours the reaction is stopped by the
addition of 75 ml aqueous saturated sodium hydrogen carbonate
solution, whereupon the product crystallizes. The dried product

is dissolved in methylene chloride and methanol and is crystallized by the addition of ligroine (b.p. 40 - 65), giving crude budesonide.

The crude budesonide product is recrystallized from methanol/water giving pure budesonide with isomer ratio A:B ≈ 1:1 (HPLC), [α]²⁵ 100.0° (c = 0.2; CH₂CL₂); M⁺ 430 (theor. 430.5)

Claims

1. Process for the manufacture of (22 R,S)-16α,17α-butylidene-dioxy-11β,21-dihydroxypregna-1,4-diene-3,20-dione

5

15

10

by reacting 11 β , 16 α , 17 α , 21-tetrahydroxypregna-1, 4-diene-3, 20-dione

20

25

30

with butanal, CH3CH2CH2CH0 in a solvent medium in the presence of a catalyst, c h a r a c t e r i s e d in that the reaction is performed in acetonitrile with p-toluenesulphonic acid as a catalyst.

35

2. Process according to claim 1, c h a r a c t e r i z e d in that the reaction is terminated by the addition of water and by adjustment of the pH of the reaction mixture.

3. Process according to claim 1 or 2, c h a r a c t e r i - z e d in that the crystals obtained upon termination of the reaction are filtered off, dissolved in methylene chloride and methanol and are then crystallized by the addition a suitable hydrocarbon, such as ligroine, hexane, cyclohexane or heptane, giving a crude product, which is then recrystallized in methanol/water to give pure budesonide.

INTERNATIONAL SEARCH REPORT

International Application No PCT/SE 90/00619

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) 6									
According to International Patent Classification (IPC) or to both National Classification and IPC									
IPC5: 0	07 J	71/00	· 						
II. FIELDS SEARCHED									
Minimum Documentation Searched'									
Classification System Classification Symbols									
:									
IPC5		C 07 J							
1100	Downstation Searched other than Minimum Documentation								
to the Extent that such Documents are included in Fields Searched ⁸									
				·					
SE,DK,	FI,NO d	classes as above							
		ONSIDERED TO BE RELEVANT®		`					
Category *	Citat	ion of Document, ¹¹ with indication, where appr	opriate, of the relevant passages 12	Relevant to Claim No.18					
Р,Х		4925933 (EDIB JAKUPOVIC E		1-3					
,,,,	15	May 1990.							
	Se	ee the whole document							
х	Chemic	cal Abstracts, volume 106,	no. 9, 2 March 1987,	1-3					
	((Columbus, Ohio, US), see pa	ige 641, abstract						
	6	7573q, ES, A, 543211 (Proc reparation of budesonide)	16 February 1986						
) P.	reparación de bocasantes,	,						
		OLCOGO COLTH MATHIESON CH	IEMT CAI	1-3					
Х	GB, A	, 916996 (OLIN MATHIESON CH DRPORATION) 30 January 1963), ·						
	Se	ee especially page 1, lines	45-66						
			•						
	ED A	I, 0164636 (SICOR SOCIETA	TALIANA	1-3					
A	CC	ORTICOSTEROIDI S.P.A.)	•						
	18	December 1985,							
	S	ee the whole document							
		a	•						
				The leavestant diles dete					
* Speci	al categor	ies of cited documents: 10	"T" later document published after or priority date and not in con- cited to understand the principal	r the international litting bate flict with the application but ple or theory underlying the					
A doc	sidered to	ning the general state of the art which is not be of particular relevance ent but published on or after the international	Invention						
"E" (B)	ng date	ent but published on or after the international	"X" document of particular releval cannot be considered novel or involve an inventive step	•					
"L" dod	cument whi ich is cited stion or oth	ch may throw doubts on priority claim(s) or to establish the publication date of another er special reason (as specified)	"Y" document of particular releva- cannot be considered to involve	nce, the claimed invention we an inventive step when the					
"O" document referring to an oral disclosure, use, exhibition or ments, such combination being obvious to a person skill									
other means "P" document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family									
IV. CERTIFICATION									
1		mpletion of the International Search							
11th D	ecembe	r 1990	195	8 -12- 2 1					
Internation	nal Searchi	ng Authority	Signature of Authorized Officer						
Gran Galler									
	SWE	DISH PATENT OFFICE	Anna Sjölund /////	VV 3/4 00 11 10 0					
Form PCT/ISA/210 (second sheet) (January 1985)									

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.PCT/SE 90/00619

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the Swedish Patent Office EDP file on 90-11-01. The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
US-A-	4925933	90-05-15	EP-A- JP-A-	0262108 63093795	88-03-30 88-04-25
GB-A-	916996	63-01-30	NONE		
EP-A1-	0164636	85-12-18	JP-A- US-A- US-A-	61040299 4695625 4835145	86-02-26 87-09-22 89-05-30

î.

r